

Table 3G
2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Metro Area

Page: 1
Run Date: 1 / 7 / 08

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Metro Area	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
Dane Cty	1,858	38.6	21.4	28.5	0.5	7.6	3.3
Mil. Cty	9,526	17.5	49.9	26.5	0.4	3.6	2.1
Racine	1,404	40.0	28.1	29.0	0.2	0.6	2.1
Unknown Metro Areas	18,646	71.7	3.5	14.4	3.1	4.5	2.8
Wisconsin	31,434	51.9	19.7	19.5	2.0	4.2	2.6
Nation (Prior Year)	857,244	48.9	23.5	20.0	1.3	2.4	3.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4G
2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Metro Area

Comparison of Age Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dane Cty	1,858	0.4	6.5	11.6	61.7	18.6	1.1	0.1
Mil. Cty	9,526	0.7	9.4	14.1	59.0	15.7	1.1	0.0
Racine	1,404	0.5	7.5	15.6	60.7	14.4	1.3	0.0
Unknown Metro Areas	18,646	0.2	5.4	14.5	63.8	14.9	1.1	0.0
Wisconsin	31,434	0.4	6.8	14.3	62.1	15.3	1.1	0.0
Nation (Prior Year)	857,244	0.4	6.6	13.0	60.4	18.2	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5G
2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by Metro Area

Comparison of Maternal Health Indicators (2, 3)												
Metro Area	Prepregnancy BMI (4)				Weight Gain (5)				Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight	< Ideal		> Ideal		3rd Trimester		Postpartum		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)		
Dane Cty	1,828	11.9 (4)	42.5 (1)	1,137	27.2 (3)	42.3 (1)	346	27.2 (3)	737	21.8 (2)		
Mil. Cty	9,436	9.0 (1)	50.1 (4)	6,741	27.7 (4)	42.8 (2)	1,473	37.9 (4)	4,251	34.5 (4)		
Racine	1,386	10.3 (2)	48.3 (3)	1,199	26.7 (2)	46.6 (4)	221	25.8 (2)	785	20.8 (1)		
Unknown Metro Areas	18,465	10.7 (3)	44.5 (2)	12,850	24.0 (1)	45.7 (3)	2,809	25.0 (1)	7,623	21.8 (2)		
Wisconsin	31,115	10.3	46.2	21,927	25.4	44.7	4,849	29.1	13,396	25.8		
Nation (Prior Year)	791,556	11.3	44.2	667,778	25.8	43.4	114,910	31.2	279,621	30.1		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6G
2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by Metro Area

Comparison of Maternal Behavioral Indicators ^(2, 3)						
Metro Area	Medical Care				WIC Enrollment	
	1st Trimester		No Care		1st Trimester	
	Number	% (Rank)	% (Rank)		Number	% (Rank)
Dane Cty	751	83.6 (2)	0.1 (3)		1,669	22.9 (4)
Mil. Cty	3,369	77.6 (4)	0.1 (3)		8,324	30.0 (1)
Racine	387	79.6 (3)	0.0 (1)		1,245	27.2 (3)
Unknown Metro Areas	7,514	86.9 (1)	0.0 (1)		16,568	29.9 (2)
Wisconsin	12,021	83.9	0.0		27,806	29.4
Nation (Prior Year)	498,811	77.6	8.2		722,757	32.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7G
2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Smoking Indicators by Metro Area

Comparison of Smoking Indicators (2, 3)										
Metro Area	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)		Number	% (Rank)	Number	% (Rank)	
Dane Cty	1,218	22.8 (2)	1,650	16.2 (2)	240	26.7 (3)	842	16.6 (1)	1,233	8.7 (2)
Mil. Cty	6,498	21.6 (1)	8,700	13.1 (1)	1,231	31.4 (1)	4,481	25.4 (3)	6,481	14.1 (4)
Racine	885	31.6 (3)	1,272	21.6 (3)	252	27.0 (2)	580	24.3 (2)	922	7.5 (1)
Unknown Metro Areas	12,471	40.0 (4)	16,972	25.7 (4)	4,492	25.2 (4)	8,398	25.9 (4)	12,404	10.0 (3)
Wisconsin	21,072	33.0	28,594	21.1	6,215	26.5	14,301	25.1	21,040	11.0
Nation (Prior Year)	677,683	30.0	553,373	19.2	140,038	33.0	362,509	28.5	443,894	26.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8G
2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by Metro Area

Comparison of Infant Health Indicators ^(2, 3)									
Metro Area	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	1,163	7.7 (1)	8.1 (3)	1,415	9.5 (1)	983	3.0 (3)	1,154	77.3 (1)
Mil. Cty	5,773	10.2 (4)	6.5 (1)	7,019	12.3 (4)	4,560	3.6 (4)	5,929	59.9 (3)
Racine	899	8.1 (3)	7.2 (2)	1,084	10.4 (3)	743	2.7 (1)	900	57.4 (4)
Unknown Metro Areas	11,821	8.0 (2)	8.5 (4)	14,396	10.3 (2)	9,775	2.8 (2)	11,863	65.0 (2)
Wisconsin	19,656	8.7	7.9	23,914	10.8	16,061	3.1	19,846	63.9
Nation (Prior Year)	707,018	8.3	7.0	636,104	12.4	583,078	3.8	704,334	59.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 3G
2004 - 2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Metro Area
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾							
Metro Area	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other/Unknown %
Dane Cty	5,587	42.2	21.2	26.8	0.3	8.1	1.4
Mil. Cty	27,866	19.4	50.7	24.6	0.5	3.7	1.2
Racine	3,567	42.2	27.0	29.0	0.2	0.4	1.1
Unknown Metro Areas	56,331	74.8	3.5	12.7	3.3	4.4	1.3
Wisconsin	93,351	55.1	19.5	17.7	2.1	4.3	1.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4G
2004 - 2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Metro Area
Combined 3 Years

Page: 8
Run Date: 1 / 7 / 08

Comparison of Age Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dane Cty	5,587	0.3	5.6	12.2	61.9	18.8	1.3	0.0
Mil. Cty	27,866	0.8	9.2	14.1	58.9	15.8	1.2	0.0
Racine	3,567	0.7	7.9	14.4	61.2	14.8	1.1	0.0
Unknown Metro Areas	56,331	0.3	5.5	14.7	63.1	15.3	1.1	0.0
Wisconsin	93,351	0.4	6.7	14.3	61.7	15.7	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5G
2004 - 2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by Metro Area
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)										
Metro Area	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	4,609	11.7 (4)	40.5 (1)	3,330	32.3 (3)	38.0 (2)	1,035	30.9 (2)	2,811	23.4 (3)
Mil. Cty	26,322	9.1 (1)	48.7 (4)	21,773	26.8 (2)	41.3 (3)	4,352	40.4 (4)	13,862	33.6 (4)
Racine	2,926	10.8 (2)	45.3 (3)	2,627	47.0 (4)	31.9 (1)	555	31.9 (3)	1,993	22.6 (2)
Unknown Metro Areas	51,509	11.3 (3)	43.0 (2)	41,915	25.0 (1)	42.6 (4)	8,439	26.1 (1)	24,692	21.8 (1)
Wisconsin	85,366	10.7	44.7	69,645	26.8	41.6	14,381	31.0	43,358	25.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).

(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6G
2004 - 2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by Metro Area
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)						
Metro Area	Medical Care				WIC Enrollment	
	1st Trimester		No Care		1st Trimester	
	Number	% (Rank)	% (Rank)		Number	% (Rank)
Dane Cty	3,653	74.8 (2)	12.3 (3)		5,225	25.0 (4)
Mil. Cty	19,103	72.4 (3)	10.6 (2)		26,314	32.3 (3)
Racine	2,176	60.7 (4)	26.4 (4)		3,248	34.6 (1)
Unknown Metro Areas	39,817	79.3 (1)	9.8 (1)		52,404	33.2 (2)
Wisconsin	64,749	76.4	10.7		87,191	32.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 7G
2004 - 2006 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by Metro Area
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Metro Area	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	3,725	29.4 (2)	4,417	17.8 (2)	832	34.3 (1)	3,299	30.9 (2)	4,339	25.8 (2)
Mil. Cty	19,888	25.0 (1)	23,818	15.7 (1)	4,238	30.9 (2)	17,988	30.6 (1)	23,413	25.6 (1)
Racine	2,449	34.7 (3)	2,892	23.2 (3)	705	26.8 (4)	2,167	35.1 (3)	2,915	25.9 (3)
Unknown Metro Areas	39,481	45.5 (4)	48,718	27.9 (4)	15,733	28.0 (3)	35,556	41.0 (4)	47,054	32.6 (4)
Wisconsin	65,543	37.9	79,845	23.5	21,508	28.8	59,010	37.0	77,721	29.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8G
2004 - 2006 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by Metro Area
Combined 3 Years

Page: 12
Run Date: 1 / 7 / 08

Comparison of Infant Health Indicators ^(2, 3)									
Metro Area	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	4,168	7.4 (2)	8.0 (2)	4,390	10.1 (2)	3,610	3.0 (2)	4,190	74.3 (1)
Mil. Cty	21,677	9.5 (4)	6.8 (1)	23,001	12.7 (4)	18,317	3.6 (4)	21,793	59.6 (3)
Racine	2,533	7.5 (3)	8.8 (3)	2,629	9.7 (1)	2,128	3.3 (3)	2,484	57.1 (4)
Unknown Metro Areas	44,068	7.3 (1)	8.9 (4)	45,817	10.7 (3)	37,338	2.7 (1)	44,217	63.4 (2)
Wisconsin	72,446	8.0	8.2	75,837	11.3	61,393	3.0	72,684	62.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.